



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

The author next relates his experiments and observations upon the urine and urinary organs of lizards. He examined four species, —the gecko, iguana, the kobbera-guion, (described by Knox,) and the alligator. The kidneys vary in size; each ureter has a papilla situated in the receptacle; in other respects the structure resembles that of snakes. The secretion is also nearly similar; that of the alligator contains, besides uric acid, carbonate and phosphate of lime; in one case it smelt strongly like musk.

In two species of the testudo, Dr. Davy found the kidneys lobulated like those of the preceding animals. In the bladder both of the turtle and tortoise he found flakes of uric acid in a transparent liquid, containing mucus and common salt, but no urea.

On a Mal-conformation of the Uterine System in Women; and on some Physiological Conclusions to be derived from it. In a Letter to Sir Everard Home, Bart. V.P.R.S. from A. B. Granville, M.D. F.R.S. F.L.S. Physician in ordinary to H. R. H. the Duke of Clarence; Member of the Royal College of Physicians, and Physician-Accoucheur to the Westminster General Dispensary. Read April 16, 1818. [Phil. Trans. 1818, p. 308.]

The uterus described in this paper had acquired its full development upon the right side only. The left side exhibited a straight line, about half an inch distant from its centre. Upon this side also all the appendages of the uterus were deficient, though their rudiments might be traced. This woman was the mother of eleven children of both sexes, and had been delivered of twins, male and female, a few days before her death, which was occasioned by diseased heart and aneurism of the aorta.

Dr. Granville remarks that this is the first case upon record which disproves the opinion that the different sides of the uterus are concerned in the production of the two sexes. It also shows that twins of both sexes may be derived from one ovarium.

This paper concludes with some remarks upon supposed cases of superfœtation.

New Experiments on some of the Combinations of Phosphorus. By Sir H. Davy, LL.D. F.R.S. V.P.R.I. Read April 9, 1818. [Phil. Trans. 1818, p. 316.]

Since the author's former communication upon the above subject to the Royal Society, various researches have been brought forward, differing in their results from his own as well as from each other. Sir Humphry concluded that the phosphoric acid contained about three fifths its weight of oxygen, or twice that contained in the phosphorous acid. Berzelius considers the phosphoric acid as composed of 100 phosphorus + 128.17 oxygen; and Dulong, of 100 phosphorus + 124.5 oxygen: and both these chemists consider the